



# Ohio Immunization Summary for School Attendance

VACCINES	FALL 2021 IMMUNIZATIONS FOR SCHOOL ATTENDANCE
<b>DTaP/DT Tdap/Td</b> Diphtheria, Tetanus, Pertussis	<p><b>Kindergarten</b> Four or more doses of DTaP or DT, or any combination. If all four doses were given before the fourth birthday, a fifth dose is required. If the fourth dose was administered at least six months after the third dose, and on or after the fourth birthday, a fifth dose is not required.*</p> <p><b>Grades 1-12</b> Four or more doses of DTaP or DT, or any combination. Three doses of Td or a combination of Td and Tdap is the minimum acceptable for children ages 7 and older, with proper spacing.</p> <p><b>Grades 7-12</b> One dose of Tdap vaccine must be administered on or after the 10<sup>th</sup> birthday. **</p>
<b>POLIO</b>	<p><b>K-11</b> Three or more doses of IPV. The final dose must be administered on or after the fourth birthday regardless of the number of previous doses. If a combination of OPV and IPV was received, four doses of either vaccine are required.</p> <p><b>Grade 12</b> Three or more doses of IPV or OPV. If the third dose of either series was received prior to the fourth birthday, a fourth dose is required. If a combination of OPV and IPV was received, four doses of either vaccine are required.</p>
<b>MMR</b> Measles, Mumps, Rubella	<p><b>K-12</b> Two doses of MMR. The first dose must be administered on or after the first birthday. The second dose must be administered at least 28 days after the first dose.</p>
<b>HEP B</b> Hepatitis B	<p><b>K-12</b> Three doses of Hepatitis B. The second dose must be administered at least 28 days after the first dose. The third dose must be given at least 16 weeks after the first dose and at least eight weeks after the second dose. The last dose in the series (third or fourth dose) must not be administered before age 24 weeks.</p>
<b>VARICELLA</b> (Chickenpox)	<p><b>K-11</b> Two doses of varicella vaccine must be administered prior to entry. The first dose must be administered on or after the first birthday. The second dose should be administered at least three months after the first dose; however, if the second dose is administered at least 28 days after the first dose, it is considered valid.</p> <p><b>Grade 12</b> One dose of varicella vaccine must be administered on or after the first birthday.</p>
<b>MCV4</b> Meningococcal	<p><b>Grades 7-11</b> One dose of meningococcal (serogroup A, C, W, and Y) vaccine must be administered prior to entry.</p> <p><b>Grade 12</b> Two doses of meningococcal (serogroup A, C, W, and Y) vaccine must be administered prior to entry. ****</p>

**NOTES:**

- Vaccine should be administered according to the most recent version of the *Recommended Immunization Schedule for Children and Adolescents Aged 18 Years or Younger* or the *Catch-up Immunization Schedule for Persons Aged 4 Months Through 18 Years Who Start Late or Who Are More Than 1 Month Behind*, as published by the Centers for Disease Control and Prevention's Advisory Committee on Immunization Practices. Schedules are available for print or download at <https://www.cdc.gov/vaccines/schedules/index.html>.
- Vaccine doses administered less than or equal to four days before the minimum interval or age are valid (grace period). Doses administered greater than or equal to five days earlier than the minimum interval or age are not valid doses and should be repeated when age appropriate. If MMR and Varicella are **not** given on the same day, the doses **must** be separated by at least 28 days with no grace period.
- For additional information, please refer to the [Ohio Revised Code 3313.67](#) and [3313.671](#) for School Attendance and the ODH Director's Journal Entry (available at <http://www.odh.ohio.gov>, [Immunization: Required Vaccines for Childcare and School](#)). These documents list required and recommended immunizations and indicate exemptions to immunizations.
- **Please contact the Ohio Department of Health Immunization Program at (800) 282-0546 or (614) 466-4643 with questions or concerns.**

\* Recommended DTaP or DT minimum intervals for kindergarten students are four weeks between the first and second doses, and the second and third doses; and six months between the third and fourth doses and the fourth and fifth doses. If a fifth dose is administered prior to the fourth birthday, a sixth dose is recommended but not required.

\*\* Tdap can be given regardless of the interval since the last tetanus or diphtheria-toxoid containing vaccine. Children age 7 or older with an incomplete history of DTaP should be given Tdap as the first dose in the catch-up series. If the series began at age 7-9, the fourth dose must be a Tdap given at age 11-12 years. If the third dose of Tdap is given at age 10 years, no additional dose is needed at age 11-12 years.

\*\*\* The final polio dose in the IPV series must be administered at age 4 or older with at least six months between the final and previous dose.

\*\*\*\* Recommended MCV4 minimum interval of at least eight weeks between the first and second doses. If the first dose of MCV4 was administered on or after the 16<sup>th</sup> birthday, a second dose is not required. If a pupil is in 12<sup>th</sup> grade and is 15 years old or younger, only one dose is required. Currently, there are no school entry requirements for meningococcal B vaccine.